

Kibkhd bkihevah\ZgguoRbferlemcgs b dh\

:\_jabg : : <eZ^bfbjh\ < E >hjhr\_gdh G < J\_amevIZlu Z\bZmq\_lguo  
^\_egby b qbke\_gghklb dblhh[jZaguo \ Ho-hikdhf] fhk\_ \ \ GZmqgh  
bkke^h\ZI\_e-vkdb ihZ hjkdbf fe\_dhiblZxsbf \ k\_\jghc qZklb Lbojh]h hd\_Zg  
F >%HUJLQ \$ \$ 9ODGLPLURY 9 Results of Airborne Research on Distribution and Abundance  
of Cetaceans in the Sea of Okhotsk in 1985. Pp. 128 in Research work on marine mammals in the  
North Pacific in 1984-1985. Moscow

:\_jabg : : <eZ^bfbjh\ < E >hjhr\_gdh G < J\_amevIZlu Z\bZmq\_lguo  
qbke\_gghklb iheyjguo k\_juo dblh\ b [\_emo-b \ H]h Ekavhf LfB G\_JH -60  
[Berzin A.A., Vladimirov V.L., Doroshenko N.V. 1990 Results of airborne research on distribution and abundance  
of bowhead, gray and beluga whales in the Sea of Okhotsk in 1985. Proceedings of TINRO, vol. 112:  
51-60]

Q\_ebgp = 2010. F\_lh^bjz k\_q\_d qbke\_gghklb Delphinapterus leucas) ih^Zgguf \u[hjhqgh]h  
609615 \ Fhjkdfe\_dhiblZxsb\_eZjdlb kfhjgbd gZmqguo ljm^h\ DZebgbg]jZ^ > & K  
2010c. The method of beluga (Delphinapterus leucas) number estimation on data of aerial surveys  
Marine Mammals of the Holarctic. Collection of Scientific Papers. Kaliningrad

Q\_ebgp \ G. = e]hbjlf jZkq\_lZ qbke\_gghklb Delphinapterus leucas ih^Zgguf \u[hjhqgh]h  
Z\bZmq\_gZkk[hjg b] Chelintsev N.G. 2012. Algorithm of white whales (Delphinapterus leucas) number  
estimation based on sample aerial survey data. This book]

Q\_jghh B, <Zkbev\_G, F\_e\_glv<\_A, =eZah F. 2008. Hiul bkihevah\Zgby ^ey v b gkljmf\_gl  
guo Z\bZmq\_lh\ fhjkdbo fe\_dhiblZkrsjzd hjZ bh E < [ Z FZI\_jbZeu f\_`^mgZjh^ghc  
dhgn\_j\_gpbb © Fhjkd\_b fe\_dhiblZxsb\_ = heZjd 1971 Chelintsev N.G., Kuznetsova V.K., Vasilyuk A.J.,  
Melentyev V., Glazov D., 2008. Laboratory aircraft L410 application for instrumental aerial surveys of marine  
mammals. (In the G L W L R Q 3 0 D W H U L D O V R I W K H , Q W H U Q D W L R Q D O & R Q I H U H  
Odessa). Pp. 132-137]

= eZah F., RiZdH<, Dmag\_p.F, Z\bZg h B, FmoZf\_ErF, Jh`gh<  
lj\_ ^ \ Z j b l \_ e v g w \_ h ] b j h k e \_ ` b \ Z g b y j \_ f \_ s \_ g b [ c \_ e m o  
(Delphinapterus leucas \ ;\_ehf fhj\_ \ - ]]  
Bgklblml ijh[e\_f wdheh]bb b w\hexpbb bf : G K\_\\_jph\Z J:G Fhk d\Z Jhkkb

Glazov D.M, Shpak O.V., Kuznetsova D.M., Ivanov D.I., Mukhametov L.M., Rozhnov V.V.  
**Preliminary results of tracking the beluga whale (Delphinapterus leucas) movements in the White Sea in 2010-2011**  
A.N. Severtsov Institute of Ecology and Evolution RAS, Moscow, Russia

:\_em Delphinapterus leucas y\ey\_lky ^\_bg  
\b^hf dblhh[jZaguo ihklhyggh h[b  
fhj\_ H[sZy \_\_ qbke\_gghklv ih  
Z\bZmq\_lh\ b h]p\_gb\Z\_lky hl  
hkh[\_c [\_a mq\_lZ `b\hlguo ih^  
^j Z \*OD]RY HW DO  
jbh^ qZklv ihimeypbb g\_ f\_g\_\_  
:\_ehf fhj\_ =eZah\ b ^j [ JZg  
qlh h^g ZhevnyZ qZklv [\_ehfhjkd bgoZ [\v  
abfgbc i\_jbh^ fb]bjm\_l \ ;Zj\_gp\_\h

The beluga whale (Delphinapterus leucas) is the only cetacean species that permanently resides in the White Sea. Based on summer aerial surveys in 2008 and 2010, the total abundance is estimated to be 5000 to 7500 individuals without correction for belugas underwater. (eZah F. 2006, 2008, 2010; Glazov et al. 2007). In winter, a considerable part of population, at least 2000 individuals, remains in White Sea water area (eZah F. 2010). Until now it has been suggested that in winter a big

Z ^jm]Zy f\_gvrZy h[bIZ\_l \ ;\_ehf  
l\_q\_gb\_ \k\_]h ]h^Z Hk\_ ^euc h[jZa  
©kZfhqvya qZklv [\_ehfhjkdhc ihime  
kIZ^Z h\leuo kZfph\ fb]jbjmxl gZ [h  
gby \ lhf qbke\_ \hafh`gh b aZ ij\_  
;\_evdh\bq KIZ^Z kZfhd  
]h^Z \ ]h^ h[bIZxl e\_lhf \ g\_kdhev  
jZchgZ@f\_kIZo j\_ijh^mdlb\guo kdn  
dh\bq Q\_jg\_pdbc b ^j  
^hklh\\_jguo ^Zgguo ih abfg\_fm jZk  
g\_l

lhiuldZ bamq\_gby i\_j\_f\_s\_gbc b kl  
\Zgby [\_emoZfb Zd\Zlhjbb ;\_eh]hZ  
eZkv lhevdh \ ] K\\_lhq\_\ l b g  
gud\_j\_ ^Zlqbd [ue mklZgh\ e\_g gZ dk  
hg\_ i Ehir\_gv]Z >\bgkdbc aZeb\  
bxgy ] ih fZjIZ ] <k\_v  
g\_ ihdb^ZeZ Zd\Zlhjbb ;\_eh]h fhj  
>\bgkdhfm Hg\_`kdhfm aZeb\Zf b  
;Zk\_cgm fhjy

P\_evx gZr\_c jZ[hlu [ueh ijhke\_^blv  
i\_j\_f\_s\_gby \ajhkeuo [\_emo \ abfgb

F\_q\_gb\_ [\_emo ijh\h^beb \ hdly[j\_  
mklv\_ j <Zjam]Z FmjfZgkdZy h[hZ  
^beky f\_lh^hf aZf\_IZ k\_lb k f\hhejg  
j\_]Z \ gb`g\_f l\_q\_gbb j\_db Fbr\_g  
[hevr\_b\_]jmiiu ba g\_kdhevdko\_jh]em  
[\_eh]h p\\_IZ aZoh^b\rb\_ \ j\_dm  
\ke\_^ aZ ]hj[mr\_c b k\_f]hc <k\_ ij  
gb\_ iheZ fhjnhf\_ljbq\_kdb\_ ijhf\_j`b  
^ey ]\_g\_l bq\_kdbo bkke\_ ^h\Zgbc b  
\uo f\_lhd ijh\h^bebkv \ \h^\_ m [h  
eh kf >ey hlike\_`b\Zgby i\_j\_f\_  
bkihevah\Zeb kimlgbdh\ u\_ i\_j\_ ^Zb  
kl\_fu \$UJRV ijhba\h^k\IZkA: Hh@kkl  
lhju\_ dj\_ibeb ijb ihfhsb ijbiZygguo  
ba g\_j`Z\\_xs\_c klZeb d lj\_f g\_ ceh  
ff ijhims\_gguf q\_j\_a ih^dh`guc \_  
emo i\_j\_ ^ ^hjkZevguf ]j\_[g\_f

<k\_]h [ueh hleh\ e\_gh kZfph\ \  
] gZ \k\_o `b\blgrablZgbe\_e\_gu  
qbdb lZ[e <klyab k g\_[hevrhc i  
jZ[hlu ^\mo i\_j\_ ^Zlqbdh\ b ^  
[ueb bkdexq\_gu ba ZgZebaZ

l\_j\ u\_ f\_kypu hgdZqZeh ^\_dZ[jy d  
ly[jv ±fhg\_p yg\Zjy ] i hkt  
i\_j\_ ^Zlqbdh\ [\_emob ^\_j`Zebkv e h  
\Z kh\\_jrZy g\_agZqbl\_evgu\_ \ df  
i\_j\_f\_s\_gby [\_emob hleh\Z  
hleh\Z ] \^hev Dhevkdh]h ih[  
\]em[v DZg^ZeZdrkdh]h aZeb\Zleat

portion of the White Sea beluga migrates to the E  
ents and Kara seas, while the smaller ones  
the White Sea throughout a year. Residential are  
the females in the White Sea populations; the a  
male groups migrate long distances, probably,  
yond the White Sea water area (evdh 1995,  
2004). In summer female units inhabit several  
SHUPDQH QV UHJLRQV WKH  
JUHJDW;L QdH`1995, 2004, Q\_jg\_pdb  
^j 2002). As for beluga distribution in winter per  
od, the reliable information is missing.

An attempt to study beluga movements and th  
strategy for habitat use in the White Sea has be  
undertaken only once, in 2005K(\\_lhqb^aj  
2007). A single transmitter was deployed on a bel  
female in the region of Lopshenga village (Dvinsk  
Bay) and transmitted from June 26 2005 to March  
2006. During this period, the beluga had not left th  
water area of the White Sea and moved betw  
Dvinskoy, Onezhsky Bays and the central parts (E  
sin) of the sea.

Our study aimed at tracking the adult beluga mo  
vements in the winter period.

Satellite tagging was conducted in October 2010 an  
2011 in the Varzuga river mouth (Murmansk regio  
Belugas were captured by a net thrown from a r  
tor-boat along the coast in the lower part of the riv  
Small groups of light grey and white belugas th  
followed pink and Atlantic salmon running up the  
river during the high tide were targeted. All pec  
dures (sex definition, morphometric measureme  
skin biopsy sampling for genetic analysis and  
deployment) were conducted near the shore at  
depth of approx. 30cm. For beluga movement trac  
ing we used Argos V\ V W H P V D W H O O L V  
V D U ` = \$ - 2 \$ 3 6 5 X V V L D 7 K H  
cables of the tags were attached to 3 nylon r  
' PP WKDW ZHUH V X E G H U P I  
back of a beluga whale in front of the dam bridge.

In total, 5 males were captured in 2010 and 3 mal  
in 2011; all of them were tagged (Table). Due to  
short period of transmission in 2011 (21 and 3 da  
two tags were excluded from analysis.

First months after tag deployment (October  
December 2010, and October 2011 to January  
2012), belugas either kept near the place of cap  
performing local movements, within 700km,  
along The Kola coast into the Gorlo or Kandalak  
sky Bay (3 belugas captured in 2010, and 1 bel  
captured in 2011), or travelled long distances (up  
300 km) from the Varzuga mouth to the Gorlo of t

White Sea following the coast line within a 100m isobath (2 belugas captured in 2010) (Fig. 1).

Table. The parameters of tagged belugas and duration of tag transmission.

ID	Sex	Color	Length	Transmission period		
				Start	Stop	Total days
61743	f	White	378 kf	27.10.2010	29.05.2011	215
61808	f	White	332 kf	30.10.2010	29.05.2011	212
61740	f	White	373 kf	30.10.2010	02.05.2011	185
61739	f	White	322 kf	30.10.2010	26.05.2011	209
61746	f	White	332 kf	30.10.2010	24.06.2011	241*
61745	f	White	410 kf	01.10.2011	22.10.2011	21
112083	f	Whitegray	380 kf	27.10.2011	is still ON**	>150
61742	f	Whitegray	370 kf	29.10.2011	01.11.2011	3




Fig. 1. The regions of the White Sea utilized by belugas according to satellite tracking data during the period of: A - from the end of October 2010 to mid-April 2011, B - from the end of October 2011 to mid-April 2012. Triangle marks the place of tagging. Dark-gray color indicates an area where the whales had been before the active ice formation began (see text)


In 2010, according to the National Ice Center <http://www.natice.noaa.gov> and generalized maps from <http://www.aari.ru> an active ice formation started in mid-December, and the next winter only in the end of January 2012. Along with a gradual land-floe formation, beluga movement pattern

oh^b|hl ijb[j\_`ghc ihehku b ihkl\_i  
 hklZevgmX Zd\Zlhjbx fhjy hkh[\_gg  
 qZklv b =hjeh

changed: the whales left the coastal zone and began to utilize the rest of the sea water area, particularly its central part and the Gorlo.

AZ \k\_ \j\_fy gZ[ex^\_gby [\_emob  
 [hdh \ DZg^ZeZdrkdbc aZeb\ qihuf  
 khdhc kiehq\_gghklvx ev^h\ \\_aZlfv  
 gu aZoh^u [\_emo \ f\_edh\h^guc  
 x`g\_\_ Kheh\\_ph\bo h

During the entire period of tracking, belugas never travelled far into Kandalakshsky Bay, which may be explained with a high ice density in the Bay. Also, belugas were not observed entering shallow Onezhsky Bay to the south of Solovetskie Islands.



Jbk Ko\_fZ i hk\_s\_gby [\_zmo h  
 lhjbb ;\_eh]h fhjy ih ^Zgguf kim  
 ijhke\_`b\Zgby k k\_j\_`bgu Zij\_ey  
 hdhgqZgby jZ[hlu i\_j\_`ZlqbdZ  
 hlf\_q\_gh djm`dhf gZ ko\_f\_`Lj  
 hlf\_q\_ghklh\_mklZgh\`db i\_j\_`Zlq  
 [\_emo Ekfguhc p\l h[hagZqZ\_l  
 dhljhjf gZoh^beZkv [\_emoZ \  
 ^h

Fig. 2. The regions of the White Sea utilized by the beluga 61746 according to the satellite tracking data mid April 2011 until the end of tag transmitting on June 24, 2011 (mapped as a black circle). Triangle mark the place of tagging. Dark-gray color indicates an area where the beluga whale was from May 2 to May 25, 2011.

H^bg kZf\_p [\_emob ghf\_j i\_j\_i^Zl  
 f\_s\_gby dhljh]h m^Zehkv hlike\_`  
 ] \ dhgp\_ Zij\_ey aZ ^g\_dh  
 jhgm <hjhgdb ;\_eh]h fhjy gZ ]jZg  
 fhj\_f WIZ [\_emoZ hklZgh\`beZkv  
 lhghc b\_g\_`eb k ih`Z  
 eZkv \ ij\_`eZo df hl mklvy d  
 gh\_ \\_jgmehkv \ ;Zkk\_cg ;\_eh]h  
 lZf \iehlv ^h hdhgqZgby jZ[hlu i\_j

One beluga male (tag number 61746), whose movements were tracked up to early June 2011, within days in the end of April had moved to the Voronka the White Sea on the border with the Barents. This whale stopped near the Ponoy river mouth, and during 3 weeks (May 02-25, 2011) stayed within 50km from the river mouth. Later, this individual returned to the Bassin of the White Sea and did not leave it until tag stopped transmitting (Fig..2)

I\_j\_`Zlqbd mklZgh\`e\_gguc \ Z  
 [hlZlv ^h gZklhys\_]h \j\_f\_gb k\_j\_`  
 ] [\_emoZ kh ^gy f\_q\_gby g\_ ihd  
 fhjy ijb wlhf Zdib\gh ih g\_fm i\_j  
 <

The tag deployed in 2011 is still transmitting at the time of paper submission (mid April 2012): from the day of tagging, the beluga has not left the Basin of White Sea and is actively moving in this water area (Fig. 1B).

LZdbf h[jZahf gZrb ^Zgg\_u\_ ihdZ  
 djZcg\_c f\_j\_ qZklv [\_emoZ\`khkgd  
 i\_jbh^ g\_ ihdb^Zxl Zd\Zlhjbx ;\_eh  
 gh i\_j\_f\_sZlky \\_]h ij\_`eZo V  
 lhevhd d abfmxsbf ih fg\_gbx g\_

Our data show that at least part of beluga population does not leave the White Sea during autumn-winter-spring period, but actively moves within its water area. This is valid not only for the females wintering; according to some authors (Levdh 2004), in the

;\_evdh\bq \ ;\_ehf fhj\_ kZfd  
 euf iheh\haj\_euf kZfpZf ljb wlf  
 k\yav f\_`^m gZqZehf ZdIb\guo i\_j  
 Zd\Zlhjbb h\_`fhjy b gZqZehf ZdIb\  
 jZah\Zgby \ ]h Zd\Zlhjbb  
 JZ[hIZ \uiheg\_gZ Ihklhyggh ^\_ck  
 pb\_c J:G ijb nbgZgkh\hc ih^^\_jhd  
 ]jZnbq\_kdh]h h[s\_kl\Z \ jZfdZo ljh  
 jZkijhkljZg\_gby b fb]jZpbc [\_em  
 ljm^gdh\ HHH ©Mljbrkdbc ^\_evrh  
 fhsv \hleh\\_ [\_emo

White Sea, but also for adult males. A connection between the start of active beluga movements on the White Sea water area and the beginning of active-ice formation in the sea is traced.

The study was conducted by the Permanent expedition of RAS under the Program of the beluga whale distribution and migrations, financial support by Russian Geographic Society. The authors are thankful to the employees of Utrish Dolphinarium, Ltd for assistance in beluga captures.

Kibkhd bkihevah\ZgguoRbferhncps b dh\

;\_evdh\bq < F Bkke\_`h\Zgb\_ kljmdlmju ihimeypbb [\_emo ;\_eh]h fhjy  
 bamq\_gbx b hojZg\_ fhjkdbo fe\_dhiblZxsb\_ =heZjdlbdb K[hjgbd gZm  
 ture in the White Sea beluga whales. Int. conf. on reproduction of marine mammals. M. Pp:10]  
 ;\_evdh\bq < F ;\_emoZ\_ljhi\_ckdh]h k\_\\_jZ gh\oibyc\_kk2k234^h\Zgby  
 [Belkovich V.M. 2004. Beluga whale of the European North: modern research. Fisheries 34: 32  
 =eZah\F., Q\_jghhd < B, ;helmgh G, FmoZf\_lh\F., RiZdH<, GZaZj\_gdh2006. E\_lgb\_  
 Z\bZmq\_luDelphinapterusleucas\ ;\_ehf fhj\_ \ b C. 148]h^ZoFhjkdb\_ fe  
 dhiblZxsb\_ =heZjdlbdb K[hjgbd gZm Glazov D.M., Chernook V.I., Boltunov A.N.,  
 Mukhametov L.M., Shpak O.V., Nazarenko E.A. 2006. Aerial survey of white whales (Delphinapterus leucas  
 Pall.) in the White Sea in 2000, 2002 and 2005. Pp:148 in Marine Mammals of the Holarctic. Collection of  
 Scientific Papers. SPb  
 =eZah\ > F Q\_jghhd < B @Zjbdh\ D : GZaZj\_gdh ? : FmoZf\_lh\ E F  
 [\_emDelphinapterusleucas \ bxe\_- ] \ ;\_ehf fhj\_ jZkij\_`\_e\_gb\_ b -198 ke\_gghklv  
 \ Fhjkdb\_ fe\_dhiblZxsb\_ =heZjdlbdb K[hjgbd gZm Glazov D.M., Chernook V.I.,  
 Zharikov K.A., Nazarenko E.A., Mukhametov L.M., Boltunov A.N. 2008. Aerial surveys of white whales (De  
 phinapterus leucas) in July in the White Sea (2005), distribution and abundance. Pp. -198 in Marine  
 Mammals of the Holarctic. Collection of Scientific Papers. Odessa]  
 =eZah\F., Q\_jghhd < B, GZaZj\_gdh@ZbdhD.:, RiZdH<, FmoZf\_lh\F. 2010Z E\_lg\_k jZ  
 ij\_`\_e\_gb\_ b qbke\_gDelphinapterusleucas ih blh]Zf Z\bZbkke\_`h\Zg]c Klj  
 134 \ Fhjkdb\_ fe\_dhiblZxsb\_ =heZjdlbdb K[hjgbd gZm Glazov D.M. m^h\ DZeb  
 Chernook V.I., Nazarenko E.A., Zharikov K.A., Shpak O.V., Mukhametov L.M. 2010. A Summer distribution and  
 abundance of belugas in the White Sea based on aerial survey data (2005). Pp. 134-140 in Marine mammals  
 of the Holarctic. Collection of Scientific Papers, Kaliningrad]  
 =eZah\ > F GZaZj\_gdh ? : Q\_jghhd < B B\Zgh\ > B RiZdHg< Kheh\ v  
 ghklb b hkh[\_gghklb jZkij\_Delphinapterusleucas o\ ;\_ehf fhj\_ \ fzj\_ - ] Klj  
 \ Fhjkdb\_ fe\_dhiblZxsb\_ =heZjdlbdb K[hjgbd gZm qguo ljm^h Glazov D.M., Nazarenko E.A.,  
 Chernook V.I., Ivanov D.I., Shpak O.V., Solovyev B.A. 2010. Assessment of abundance and distr  
 bution peculiarities of beluga whales (Delphinapterus leucas) in the White Sea in 2010. Pp. 140-145 in  
 Marine Mammals of the Holarctic, Collection of Scientific Papers. Kaliningrad]  
 K\\_lhq< G, K\\_lhq H G, ;hg^Zj< \, ljbs\_fbob\_gN, NjhehK; . 2007. Bamq\_gb\_ k\_ahgguo  
 fb]jZpbc [\_emDelphinapterusleucas Pall ih fhsvx ^Zlqbdh\ kimlgbdh\hc l\_e\_f\_ljbb > KL  
 fhj\_ K - \ k[ ljh[e\_fu bamq\_gby jZp bki b hojZgu ijbj\_`uk O jkh\ ;\_e  
 f\_`^mgZj dhgn k\_gly[jy ] :jo Svetlocheva N., Svetlocheva O.N., Bondarev V.A.,  
 Prischemikhin V.F., Frolov S.B. 2007. Study of seasonal migrations of belugas (Delphinapterus leucas Ball.) u  
 ing satellite telemetry sensors in the White Sea. Pp:3505 in Problems of study, sustainable use and conserv  
 tion of natural resources of the White Sea. Proceedings of the 10th International Conference. Arkhangelsk]  
 Q\_jg\_p.d.b.c;\_evdh\bq, DjZkgh\Z2002.Gh\u\_`^Zggu\_ h kljmdlmj\_ ihimeypbb [\_emo  
 fhj\_ K - \ Fhjkdb\_ fe\_dhiblZxsb\_ =heZjdlbdb F > &k\WU, Krasnova L\ \$ ' %

V.V. 2002. New data about population structure of white whales in the White Sea. Marine mammals of the Arctic, Moscow. Pp. 272-282]

Glazov D.M., Chernook V.I., Boltunov A.N., Mukhametov L.M., Shpak O.V., Nazarenko E.A. 2007. Preliminary results of the White sea beluga (*Delphinapterus leucas* Pall.) aerial surveys (2005, 2006) Abstract book of the International Workshop on Beluga Whale Research, Husbandry and Management in wild and captive environment, 9-11 March, 2007, Valencia, Spain. p.6

= h^y s\_X Z^1, ;\_e y q\_ g. d h : e l m o h. <^2, ; m j d Z g h G^2,3

**Dhimeylhjgh ih \ ^ \_ g b \_ k Z f h Eumetopias jubatus )**

GZpbhgZevguc bkke\_ ^h \ ZI\_ evkdbc KZjZih \ kdbc ] h k m ^ Z j k l \ \_ g g u c m g b  
 KZjZih \ J h k k b y  
 DZf q Z I k d b c n e b Z e L b o h h d \_ Z g k d h ] h b g k l b l m I Z ] \_ h j D Z f q Z I k d b c J : G h k l  
 3. GZpbhgZevgZy eZ[hjZihjby ih bamq\_gbx fhjkdbo fe\_dh ib l Z x s b o : e y k  
 b k k e \_ ^ h \ Z I \_ e v k d h ] h j u [ h e h \ g h ] h p \_ g l j Z G Z p b h g Z e v g h c k e m ` [ u f h j k d h  
 KR:

Godyashcheva Y.S., Belyachenko A.V.<sup>1</sup>, Altukhov A.V.<sup>2</sup>, Burkanov V.N.<sup>2,3</sup>

**Copulatory behaviour in female Steller sea lions (*Eumetopias jubatus*)**

1. Saratov State University named after N.G. Chernyshevsky, Saratov, Russia
2. Kamchatka Branch of the Pacific Institute of Geography, RAS, Petropavlovsk-Kamchatsky, Russia
3. National Marine Mammal Laboratory, Alaska Fisheries Science Center, National Marine Fisheries Service, National Oceanic and Atmospheric Administration, Seattle, USA

K b \ m E u m e t o p i a s j u b a t u s ) - d j m i g \_ c r b c i j \_ ^ k l  
 k \_ f \_ c k l \ Z m r Z k l u o l x e \_ g \_ c b f \_ x  
 \_ k l \_ k l \ \_ g g u c j \_ i j h ^ m d l b \ g u c i h l \_ g p  
 \ b ^ Z \ g \_ ^ Z \ g \_ \_ \ j \_ f y i j \_ l \_ j i \_ e Z z g  
 s \_ g b \_ i h ^ ^ \_ c k l \ b \_ f k i g u d z m j d M e r r i c k  
 et al. 1987, Loughlin et al. 1992 Burkanov and Loughlin  
 H ^ g h c b a \ Z ` g u o a Z ^ Z q \ b a r  
 ` \_ g b y q b k e \_ g g h k l b k b \ m q Z y l e y \_ h k  
 \ Z g b \_ \ h i j h k h \ ^ \_ f h ] j Z n b b b \ h k i j h  
 g m x j h e v \ j \_ r \_ g b b i j w k f Z ] b ] j Z \_ l b k k e  
 d h i m e y l h j g h ] h i h \ ^ \_ g b y k b \ m q \_ c  
 (Amos et al. 2001, Hoffman et al h l f  
 q l h m g \_ d h l h j u o \ b ^ h \ m r Z k l u o l x e  
 h j ] Z g b a Z p b b j Z a f g h ` \_ g b y a Z \ u r \_ g z  
 f \_ g l g \_ y k g h k m s b k r h \ n k Z l f h d \ h a f h b g  
 y l v g Z \ u [ h j i z j l g \_ j Z b \ d Z d h c k l  
 y l e y \_ l k y i m k d h \ u f \_ f \_ o Z g b a f h f g Z c  
 l h w l h f m h k h [ u c b g l \_ j \_ k i j \_ ^ k l Z \ e y  
 k Z f h d \ n h j f b j h \ Z g b b j \_ i j h ^ m d l b \ g  
 : e l m o h \ P \_ e v j Z [ h e l x q z e z k v \ Z  
 h k h [ \_ g g h k l \_ c d h i m e y l h j g h ] h i h \ ^ \_  
 \ \ u y k g \_ g b b \ h a f h ` g h k l b b o Z d l b \ g  
 g h ] h \ u [ h j Z  
 G Z [ e x ^ \_ g b y a Z k b \ m q Z f b i j h \ h ^ b e

Steller sea lion (*Eumetopias jubatus*) is the largest representative of the family of eared seals, with relatively low natural reproductive potential. Population of this species has recently undergone a significant decline in numbers. The causes and influencing factors are not yet clear (Merrick et al. 1987, Loughlin et al. 1992, Burkanov and Loughlin 2005). In studying the causes of decline of sea lions' population, one of the key tasks is a detailed study of demography and reproduction of this species. An important role in this task is played by study of copulatory behaviour of Steller sea lions. Many authors (Amos et al. 2001, Hoffman et al. 2003, 2007) noted that in some species of eared seals the role of males in breeding organization is overstated. At the moment it is not clear whether females are able to influence the choice of a partner, and to what extent well as what triggers the beginning of mating. Therefore, elucidating the role of females in formation of sexual relationships is of particular interest (Glazov et al. 2012). The purpose of this work was to analyse characteristics of female sea lions' copulatory behaviour and to determine whether they have a possibility of active reproductive choices.